

WHAT ARE THE BENEFITS OF **PROLASTIC?**

THE FIRST MINIMALLY INVASIVE PRODUCT TO ACHIEVE LONG-TERM RESULTS IN MEN

Injectables are usually made from resorbable materials, which reduces their effectiveness over time. Prolastic is produced from a non-resorbable, tissue-friendly material, for a long-lasting effect.

IT WILL NOT ATTACH TO THE SURROUNDING TISSUE

Prolastic will not attach to the surrounding tissue, which makes it easy to remove if so required.

IMMEDIATE EFFECT

The procedure will immediately stop any urine loss.

NO HOSPITALIZATION

The procedure is not stressful and allows you to resume your daily activities almost immediately.

SAFE

No general anesthetic and no incisions.

PROLASTIC

OUTPATIENT TREATMENT AND MINIMALLY INVASIVE

- Reliable and safe
- Long-lasting
- Fast outpatient procedure
- No urethral mucosal damage
- No migration
- No general anesthetic



PLEASE REFER TO YOUR DOCTOR FOR MORE INFORMATION

Would you like to learn more about Prolastic?
Visit www.urogynbv.com

PROLASTIC

Minimal invasive and long-lasting!



PROLASTIC

The long-term solution for male incontinence

Stress incontinence in men

Urinary incontinence can be short-term or chronic. Short-term incontinence is often caused by other health problems or treatments. Stress incontinence is a form of chronic urinary incontinence.

Stress incontinence happens when you sneeze, cough, laugh, lift objects, or do anything that puts stress or strain on your bladder and causes you to leak urine.

- Reliable and safe
- Long-lasting
- Fast outpatient procedure
- No urethral mucosal damage
- No migration
- No general anesthesia

What causes stress incontinence?

Stress incontinence often occurs following a prostatectomy or TURP. This is the (partial) removal of the prostate. After the removal of the prostate some men may experience loss of urinary control for a number of reasons. Because of the sensitive location of the prostate, its surgical removal may cause disruption of either the anatomic continence mechanism or the nerves that influence urinary control.

How is urinary incontinence in men diagnosed?

Your doctor will do a physical examination, ask questions about your symptoms and medical history, and test your urine. Often this is enough for the doctor to tell if you have stress incontinence. You may need other tests if the incontinence has several causes or if the cause is unclear.


How is it treated?

In men, incontinence is often related to prostate problems or treatments. If changes in lifestyle and pelvic floor exercises prove insufficient to control incontinence, surgery may be an option.

How can Prolastic help you?

Prolastic is an injectable bulking agent. Two components are mixed that will set after a few minutes. While the mixture is still liquid, it is injected at two positions around the urethra. This will narrow and stabilize the urethra and improve the closure mechanism.

Prolastic does not stick to the tissue and can be removed in case of overcorrection. Prolastic has a long-lasting effect since the material keeps its shape. The procedure is simple, minimally invasive and fast. The treatment lasts around 15 minutes and is performed under local anesthesia.



The material is deposited where the prostate used to be (after a radical prostatectomy). After a TURP the material will be placed close to the prostate.

Urogyn BV
Transistorweg 5a
6534 AT Nijmegen
The Netherlands

t +31(0) 24 711 41 30
info@urogynbv.com
www.urogynbv.com